

# Large-scale HI in nearby radio galaxies: segregation in neutral gas content with radio source size

B.H.C. Emonts<sup>1,2</sup>, R. Morganti<sup>1,3</sup>, T.A. Oosterloo<sup>1,3</sup>, J.M. van der Hulst<sup>1</sup>, G. van Moorsel<sup>4</sup>, and C.N. Tadhunter<sup>5</sup>.

<sup>1</sup> Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, the Netherlands

<sup>2</sup> Department of Astronomy, Columbia University, Mail Code 5246, 550 West 120th Street, New York, N.Y. 10027, USA  
e-mail: emonts@astro.columbia.edu

<sup>3</sup> Netherlands Foundation for Research in Astronomy, Postbus 2, 7990 AA Dwingeloo, the Netherlands

<sup>4</sup> National Radio Astronomy Observatory, Socorro, NM 87801, USA

<sup>5</sup> Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK

**Abstract.** We present results of a study of neutral hydrogen (HI) in a complete sample of nearby non-cluster radio galaxies. We find that radio galaxies with large amounts of extended HI ( $M_{\text{HI}} \gtrsim 10^9 M_{\odot}$ ) all have a compact radio source. The host galaxies of the more extended radio sources, all of Fanaroff & Riley type-I, do not contain these amounts of HI. We discuss several possible explanations for this segregation. The large-scale HI is mainly distributed in disk- and ring-like structures with sizes up to 190 kpc and masses up to  $2 \times 10^{10} M_{\odot}$ . The formation of these structures could be related to past merger events, although in some cases it may also be consistent with a cold-accretion scenario.

**Key words.** Galaxies: active – Galaxies: interactions – (Galaxies:) cooling flows – Galaxies: ISM – ISM: jets and outflows

## 1. Introduction

Are different types of radio sources intrinsically different or do they inhabit different galactic environments? We address this long-standing ‘nature vs. nurture’ issue in a study of the neutral hydrogen (HI) gas in nearby radio galaxies. HI gas gives insight in the content and distribution of inter-stellar medium (ISM) in the host galaxies of the radio sources. It can be traced both at large scales (in emission), as well as in the central region of the galaxy (in absorption against the radio continuum), where it may serve as fuel for the active galactic nucleus (AGN). HI can also serve as a long-lived tracer of the galaxy’s history. For example, large-scale HI structures may have formed from past merger or interaction events (e.g. Hibbard & van Gorkom 1996), which are often invoked to trigger nuclear activity.

In this Letter we present and discuss a segregation that we find in large-scale HI content between compact and extended radio sources, the latter all of Fanaroff & Riley (1974) type-I (FR-I). The results are based on a sample of 21 radio galaxies from the B2-catalogue ( $F_{408\text{MHz}} \gtrsim 0.2$  Jy) up to a redshift of  $z \approx 0.04$ . This sample is *complete*, with the restriction that we left out sources in dense cluster environments (since here large-scale gaseous features are likely wiped out on relatively short timescales) and BL-Lac objects. In addition we observed NGC 3894, which has a compact radio source with radio power com-


parable to our B2-sample sources. In total we observed 9 compact ( $< 15$  kpc) and 13 extended FR-I ( $> 15$  kpc) radio sources. The sources have a radio power  $22.0 \leq \text{Log}(P_{1.4\text{ GHz}}) \leq 24.8$  with no bias in  $P_{1.4\text{ GHz}}$  between the compact and extended sources. The radio sources are hosted by *early-type galaxies* (E and S0). Observations were made during various observing runs in the period Nov. 2002 - Feb. 2005 with the Very Large Array (VLA) in C-configuration and the Westerbork Synthesis Radio Telescope (WSRT). A full description of the sample and observing details will be presented in a future paper (Paper II).

## 2. Results

We detect HI in emission associated with 6 of our objects. The total intensity images and properties of the HI structures (if not previously published) are shown in Figure 1 and Table 1.<sup>1</sup> In most cases the HI is distributed in a fairly regularly rotating disk or ring, although a varying degree of asymmetry is still visible in these structures. For B2 0258+35, B2 0648+27 and NGC 3894 the HI structure is more than 100 kpc in size and contains at least about the HI mass of the Milky Way.<sup>2</sup> For B2 0722+30 part of the HI disk is seen in absorption against the radio continuum, therefore the total mass of the HI disk is somewhat larger than given in Table 1. The environment of B2 0722+30 (see also Fig. 1) is very dynamic and gas-rich (the

<sup>1</sup> We assume  $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$  and use the redshift (unless otherwise indicated) to calculate the HI properties.

<sup>2</sup>  $M_{\text{HI, MW}} = 5 \times 10^9 M_{\odot}$  (Henderson et al. 1982).



**Fig. 1.** 0th-moment total intensity maps of the HI emission (contours) around five nearby radio galaxies. Radio continuum is only shown for B2 1322+36 (grey contours); for the other sources, the radio continuum is unresolved (or only marginally resolved for B2 0722+30). Although HI absorption is present in all five radio galaxies, we only show the HI absorption (white contours/profile) in case it clarifies the morphology of the large-scale HI. The arrows mark the host galaxies of our sample sources, while the broken lines show the direction along which the position-velocity (PV) plots are taken. B2 0648+27: contour levels 0.22, 0.36, 0.52, 0.71, 0.95, 1.2, 1.5, 1.8,  $2.1 \times 10^{20} \text{ cm}^{-2}$  (see also Emonts et al. 2006b); B2 0258+35: levels from 0.34 to 3.0 in steps of  $0.44 \times 10^{20} \text{ cm}^{-2}$ ; B2 1322+36: 1.7, 2.3, 2.8 (black)  $\times 10^{20} \text{ cm}^{-2}$ ; absorption (white) at 17, 45, 70, 95% of peak absorption – continuum levels: from 22 to 200 in steps of  $44.5 \text{ mJy beam}^{-1}$  (grey); NGC 3894: levels 0.17, 0.49, 0.87, 1.7, 3.2, 4.6 (black)  $\times 10^{20} \text{ cm}^{-2}$ ; absorption (white) at 4, 43, 87% of peak absorption (PV: levels -1.0, -5.0, -10, -14 (grey), 1.0, 2.0, 3.0, 4.5, 6.5 (black)  $\text{mJy beam}^{-1}$ ); B2 0722+30: levels 0.67, 1.3, 1.8, 2.3, 3.0, 4.0, 5.0, 6.0, 7.0,  $8.0 \times 10^{20} \text{ cm}^{-2}$  – part of the HI disk that is observed in absorption is not plotted – (PV plot: levels -0.5, -1.4, -2.4, -3.4, -4.4 (grey), 0.5, 0.7, 0.9, 1.1, 1.3 (black)  $\text{mJy beam}^{-1}$ ).

total mass in the field of B2 0722+30 is  $M_{\text{HI}} \approx 4 \times 10^9 M_{\odot}$ . B2 1322+36 shows “blobs” of HI emission, as well as slightly extended HI absorption, in the direction of a nearby companion, possibly tracing a bridge- or tail-like structure. Although HI absorption is detected in the central beam of all the radio galaxies in Fig. 1, as well as a few extended FR-I sources in our sample (which we will present in Paper II), this Letter is restricted to the HI gas detected in *emission* in large-scale structures that stretch beyond the optical extent of the host galaxy.

The most intriguing result from this Letter is shown in Figure 2. Large amounts of extended HI (with  $M_{\text{HI}} \geq 10^9 M_{\odot}$ ) are only observed around host galaxies with a *compact* radio source, while none of the host galaxies of the more extended FR-I type radio sources shows similar amounts of large-scale HI. For the non-detections we derived a firm upper limit by estimating the total HI mass of a potential  $3\sigma$  detection smoothed across a velocity range of  $400 \text{ km s}^{-1}$  (similar to that of the detected HI disks/rings). Note that this does not exclude the possibility that the extended FR-I sources contain more conser-

**Table 1.** HI in radio galaxies. Given is the B2 name, the NGC number, the total HI mass detected in emission, the diameter of the HI structure (or distance to the host galaxy for B2 1322+36), the peak in HI surface density, and the morphology of the HI structure (D = disk, R = ring, B = “blob”).

| # | B2 Name              | NGC  | $M_{\text{HI}}$<br>( $M_{\odot}$ ) | $D_{\text{HI}}$<br>(kpc) | $\Sigma_{\text{HI}}$<br>( $M_{\odot}/\text{pc}^2$ ) | Mor.<br>HI |
|---|----------------------|------|------------------------------------|--------------------------|-----------------------------------------------------|------------|
| 1 | 0258+35              | 1167 | $1.8 \times 10^{10}$               | 160                      | 2.7                                                 | D          |
| 2 | 0648+27 <sup>a</sup> | -    | $8.5 \times 10^9$                  | 190                      | 1.7                                                 | R          |
| 3 | 0722+30              | -    | $2.3 \times 10^8$                  | 15                       | 4.1                                                 | D          |
| 4 | 1217+29 <sup>b</sup> | 4278 | $6.9 \times 10^8$                  | 37                       | -                                                   | D          |
| 5 | 1322+36              | 5141 | $6.9 \times 10^7$                  | 20                       | 3.7                                                 | B          |
| 6 | -                    | 3894 | $2.2 \times 10^9$                  | 105                      | 3.8                                                 | R          |

References: a). Emonts et al. (2006b); b). Morganti et al. (2006).

vative amounts of large-scale HI. Take for example the very nearby radio source Centaurus A, which has a total linear extent of 650 kpc. Cen A contains  $1.5 \times 10^8 M_{\odot}$  of HI in diffuse

shells outside the optical host galaxy (van Gorkom et al. 1990; Schiminovich et al. 1994). In addition, Cen A has an inner disk of  $M_{\text{HI}} \approx 4.5 \times 10^8$ , but this disk would be only marginally resolved at the distance of our sample sources and resolution of our observations and it would appear mostly in absorption against the central radio continuum.

When applying a statistical Mann-Whitney U-Test on our data, the group of compact (<15 kpc) radio sources differs in large-scale HI mass content from the group of extended (>15 kpc) radio sources already at the 95% significance level (even when leaving NGC 3894 out of the statistics). This suggests that for these nearby radio galaxies there indeed is a *segregation in large-scale neutral hydrogen content between compact and extended radio sources*.


### 3. Discussion

The question arises what could cause this segregation in HI-content between the compact radio sources and the extended FR-I radio sources? To answer this question one needs to understand more about the nature - and therefore about the origin - of these large-scale HI structures. One possibility is that they originated from merger events. In a major merger between gas-rich galaxies (e.g. Hibbard & van Gorkom 1996; Mihos & Hernquist 1996; Barnes 2002), part of the gas is transported to the central region, while another part is expelled in large structures of low surface brightness (tidal-tails, bridges, shells, etc.). If the environment is not too hostile and the gas in these large structures remains gravitationally bound to the system, it can fall back onto the galaxy and settle into a disk- or ring-like structure within a few galactic orbits (>1 Gyr; Barnes 2002). For B2 0648+27 we confirmed a merger origin through the detection of a post-starburst stellar population (Emonts et al. 2006b). Another possible formation mechanism of the large-scale HI structures is cold accretion. Simulations by Kereš et al. (2005) show that galaxies can accrete gas from the inter-galactic medium via a cold mode, i.e. the gas cools along filamentary structures without being shock-heated. On smaller scales, Kaufmann et al. (2006) show that through the cooling of hot halo gas, cold gas can be assembled onto a galactic disc.

Leaving open the exact formation mechanism, we discuss several possible explanations for the segregation in large-scale HI-content between the compact sources and the extended FR-I sources:

The first possibility is that the large-scale HI content and type of radio source both depend on the properties of the host galaxy. As we will describe in Paper II, we find that, using the available optical information, no other properties of the host galaxy correlate strongly with the presence of extended HI. However, deep optical imaging, as well as information at other wavelengths, is necessary to study in more detail the properties of the host galaxies.

The second possibility is that the radio sources heat/ionise the gas when jets/lobes propagate outward. An example of this is seen in the radio galaxy Coma A, where an HI disk (60 kpc in



**Fig. 2.** Total HI mass detected in emission plotted against the total linear extent of the radio sources. Source sizes have been taken from numerous publications (see Emonts 2006, and references therein). In case of non-detection a tight upper limit ( $3\sigma$  across  $400 \text{ km s}^{-1}$ ) is given. The numbers correspond to the sources as they are given in Table 1.

diameter) is partly ionised by radio jets/lobes expanding into it (Morganti et al. 2002). Simulations of powerful radio sources by Bicknell & Sutherland (2006) show that propagating radio jets can create a quasi-spherical, high pressured bubble of radio plasma, that drives radiative shocks in every direction through the host galaxy's ISM. This mechanism - if important also for FR-I sources - might explain how extended neutral gas structures (such as the ones in Fig. 1) may have been ionised by the extended radio sources in our sample. Alternatively, Best et al. (2006) argue that moderately powerful radio sources can self-regulate the balance between cooling and heating of hot gas surrounding these systems. In light of the cold accretion scenario as a possible formation mechanism of large-scale HI structures, this suggests that propagating radio jets may prohibit HI structures from forming in the first place.

Extensive emission-line studies (Baum et al. 1988; Baum & Heckman 1989; Baum et al. 1992) show that large-scale features of warm gas are common in nearby powerful radio galaxies, but for FR-I sources they are not as extended, massive and regularly rotating as the HI disks/rings that we find around some compact sources. Therefore, we do not expect that the extended sources in our sample have converted similar HI structures into warm emission-line gas. On the other hand, X-ray studies are necessary to investigate the presence of hot gas around our sample sources.

The third possibility is that the HI-rich compact radio sources do not grow into extended sources, either because they are frustrated by the ISM in the central region of the galaxy, or because the fuelling stops before the sources can expand. If our HI-rich radio galaxies formed through a major merger of gas-rich galaxies, part of the gas is expelled in large-scale tidal structures, while another part is transported into the central kpc-scale region (e.g. Barnes 2002). The latter could be responsible for frustrating the radio jets if they are not too powerful. Alternatively, while the geometry and the

conditions of the encounters appear to be able to form the observed large-scale HI structures, they might not be efficient in channelling substantial amounts of gas to the very inner pc-scale region. This might prevent stable fuelling of the AGN and hence large-scale radio structures do not develop.

The four most compact and most HI-rich radio sources in our sample have been studied in detail by Giroletti et al. (2005a,b) and Taylor et al. (1998). Interaction with the ambient medium has been suggested for all these sources. Although it is not clear how much gas is needed to confine a radio source (see e.g. the discussion by Holt et al. 2003), Giroletti et al. argue that the relatively low power radio sources in NGC 4278 and B2 0648+27 can not bore through the local ISM. B2 0258+35 displays variable levels of activity, suggestive of inefficient fuelling, and is therefore not expected to grow beyond the kpc scale (Giroletti et al. 2005a).

The extended FR-I sources could be fed in another way. One possibility are cooling flows (e.g. Fabian 1994), which may create a continuous supply of gas, allowing the source to grow. Since only a very small amount of gas is necessary to fuel a radio source (e.g. van Gorkom et al. 1989), another possibility is that FR-I sources are related to dry mergers between gas-poor elliptical galaxies (e.g. Colina & de Juan 1995).

### 3.1. Relation to other studies

In a study of Ly $\alpha$  emission-line halos around radio galaxies at  $z > 2$ , van Ojik et al. (1997) detect numerous cases of strong HI absorption in the Ly $\alpha$  emission-line profiles, mainly in the smaller radio sources. They prefer the possible explanation that these small radio sources reside in dense, possibly (proto) cluster environments, where the radio source vigorously interacts with the ambient medium. It is worth noting that our nearby radio galaxies were selected *not* to lie in dense clusters. Villar-Martín et al. (2003) argue that Ly $\alpha$  halos at  $z > 2$  might be cooling flow halos that could be related (progenitors?) to large-scale HI structures at low- $z$ . A detailed comparison between low and high redshift HI-rich radio galaxies deserves further attention.

Finally, large-scale HI is also found around the compact radio source PKS 1718-649 (Veron-Cetty et al. 1995), as well as around radio-quiet early-type galaxies (e.g. van Gorkom & Schiminovich 1997; Morganti et al. 1997, 2006; Oosterloo et al. 2002; Serra et al. 2006). Despite the small number statistics and difference in observational restrictions, so far no major differences in HI characteristics (detection rate, morphology and mass) appear between samples of radio-quiet and our sample of radio-loud early-type galaxies (see Emonts et al. 2006a). This might indicate that the radio-loud phase (although not necessarily at the level of the more powerful FR-I sources) is just a short period in the lifetime of many, or maybe even all, early-type galaxies.

## 4. Conclusions

We find a segregation in large-scale neutral hydrogen content with radio source size in a complete sample of nearby non-cluster radio galaxies. Large-scale HI structures that stretch

beyond the optical host galaxy and that have  $M_{\text{HI}} \gtrsim 10^9 M_{\odot}$  detected in emission are found only around galaxies with a compact radio source, while the more extended FR-I radio galaxies do not show similar amounts of large-scale HI. If confirmed by studies of larger samples, the large-scale neutral hydrogen content may be a specific property of the host galaxy for various types of radio sources throughout the universe.

*Acknowledgements.* We would like to thank J. van Gorkom for useful discussion and the referee for the detailed comments that improved this paper. Part of this project is funded by the Netherlands Organisation for Scientific Research (NWO) under Rubicon grant 680.50.0508. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. The Westerbork Synthesis Radio Telescope is operated by the Netherlands Foundation for Research in Astronomy (ASTRON) with support from NWO.

## References

- Barnes, J. E. 2002, MNRAS, 333, 481
- Baum, S. A. & Heckman, T. 1989, ApJ, 336, 681
- Baum, S. A., Heckman, T. M., Bridle, A., van Breugel, W. J. M., & Miley, G. K. 1988, ApJS, 68, 643
- Baum, S. A., Heckman, T. M., & van Breugel, W. 1992, ApJ, 389, 208
- Best, P. N., Kaiser, C. R., Heckman, T. M., & Kauffmann, G. 2006, MNRAS, 368, L67
- Bicknell, G. V. & Sutherland, R. S. 2006, AN, 327, 235
- Colina, L. & de Juan, L. 1995, ApJ, 448, 548
- Emonts, B. H. C. 2006, Ph.D. Thesis
- Emonts, B. H. C., Morganti, R., & Oosterloo, T. A. 2006a, New Astron. Rev., in press.
- Emonts, B. H. C., Morganti, R., Tadhunter, C. N., et al. 2006b, A&A, 454, 125
- Fabian, A. C. 1994, ARA&A, 32, 277
- Fanaroff, B. L. & Riley, J. M. 1974, MNRAS, 167, 31P
- Giroletti, M., Giovannini, G., & Taylor, G. B. 2005a, A&A, 441, 89
- Giroletti, M., Taylor, G. B., & Giovannini, G. 2005b, ApJ, 622, 178
- Henderson, A. P., Jackson, P. D., & Kerr, F. J. 1982, ApJ, 263, 116
- Hibbard, J. E. & van Gorkom, J. H. 1996, AJ, 111, 655
- Holt, J., Tadhunter, C. N., & Morganti, R. 2003, MNRAS, 342, 227
- Kaufmann, T., Mayer, L., Wadsley, J., Stadel, J., & Moore, B. 2006, MNRAS, 370, 1612
- Kereš, D., Katz, N., Weinberg, D. H., & Davé, R. 2005, MNRAS, 363, 2
- Mihos, J. C. & Hernquist, L. 1996, ApJ, 464, 641
- Morganti, R., de Zeeuw, P. T., Oosterloo, T. A., et al. 2006, MNRAS, 371, 157
- Morganti, R., Oosterloo, T. A., Tinti, S., et al. 2002, A&A, 387, 830
- Morganti, R., Sadler, E. M., Oosterloo, T., Pizzella, A., & Bertola, F. 1997, AJ, 113, 937
- Oosterloo, T. A., Morganti, R., Sadler, E. M., Vergani, D., & Caldwell, N. 2002, AJ, 123, 729

- Schiminovich, D., van Gorkom, J. H., van der Hulst, J. M., & Kasow, S. 1994, *ApJ*, 423, L101
- Serra, P., Trager, S. C., van der Hulst, J. M., Oosterloo, T. A., & Morganti, R. 2006, *A&A*, 453, 493
- Taylor, G. B., Wrobel, J. M., & Vermeulen, R. C. 1998, *ApJ*, 498, 619
- van Gorkom, J. & Schiminovich, D. 1997, in *ASP Conf. Ser.* 116, 310
- van Gorkom, J. H., Knapp, G. R., Ekers, R. D., et al. 1989, *AJ*, 97, 708
- van Gorkom, J. H., van der Hulst, J. M., Haschick, A. D., & Tubbs, A. D. 1990, *AJ*, 99, 1781
- van Ojik, R., Roettgering, H. J. A., Miley, G. K., & Hunstead, R. W. 1997, *A&A*, 317, 358
- Veron-Cetty, M.-P., Woltjer, L., Ekers, R. D., & Staveley-Smith, L. 1995, *A&A*, 297, L79+
- Villar-Martín, M., Vernet, J., di Serego Alighieri, S., et al. 2003, *MNRAS*, 346, 273